

SUCCESSFUL MANAGEMENT OF FETAL DYSTOCIA DUE TO DOG SITTING POSTURE WITH NAPE PRESENTATION IN A JENNY

R. RUTHRAKUMAR*, G. SATHRIYAN, D. GOPIKRISHNAN, A. GANESAN, M. SELVARAJU¹
and K. SENTHIL KUMAR

Department of Veterinary Gynaecology and Obstetrics, ¹Dean, Veterinary College and Research Institute, Namakkal
Tamil Nadu Veterinary and Animal Sciences University (TANUVAS), Namakkal-637 002, Tamil Nadu, India

Received: 02.04.2023; Accepted: 17.06.2023

SUMMARY

Obstetrical emergencies like dystocia are considered to be the life threatening disorder of dam and foetus. The long extremities predispose the foetus to pave the way for postural abnormalities. In donkeys, the epidemiological data of occurrence and etiology of dystocia were studied not well known or is still in its infancy. The most prominent attributable factor of dystocia in donkeys were often due to postural abnormalities leading to severe injuries to reproductive tract and increased odds for mortality of foetus. A full term pregnant eight year old female donkey in her third parity was brought to Large Animal Obstetrical Unit, VCC, VCRI, Namakkal with the history of straining since two days and water bag ruptured one day before. General clinical examination revealed elevated body temperature (104 °F) with right lateral recumbency and vulvar edema. Haematology revealed leucocytosis with neutrophilia. Upon vaginal examination, the foetus was found in dog sitting posture with nape presentation and absence of pedal reflex. Hence the case was diagnosed as dystocia due to postural defects. Treatment include the thorough lubrication of the foaling passage under epidural anaesthesia followed by mutation operation and traction a dead male foetus relieved per vaginally. As a post obstetrical management the donkey was treated with parenteral fluids, antibiotics and anti-inflammatory administered for five days.

Keywords: Dog sitting posture, Dystocia, Jenny

How to cite: Ruthrakumar, R., Sathriyan, G., Gopikrishnan, D., Ganesan, A., Selvaraju, M. and Senthil Kumar K. (2023). Successful management of fetal dystocia due to dog sitting posture with nape presentation in a Jenny. *The Haryana Veterinarian* 62(2): 152-153.

Dystocia in equids is always an emergency condition because within an hour of onset of second stage of labour, placental separation from lining of the uterus begins and leads to progressive asphyxia of foal (Noakes, 2001). Jenny's reproductive characteristics are quite similar to those of a mare because they both belong to the genus *Equus*. In donkeys, the average gestation period lasts between 10½ and 14½ months. Dystocia in Jenny is very rare and occur occasionally, about 1 to 4 % of foaling. One of the most difficult obstetrical challenges is the dogsitting posture (Card, 2002). The incidence of dog-sitting posture is reported to be 0.7% (Baldwin *et al.*, 1991). However, the per-vaginal delivery of foal is difficult due to its long extremities (Arthur *et al.*, 2001). The present study deals with the successful management of fetal dystocia due to dog sitting posture with nape presentation in a jenny.

A full term pregnant eight year old female donkey in her third parity was brought to Large Animal Obstetrical Unit, VCC, VCRI, Namakkal with the history of straining since 2 days. The water bag appeared and ruptured one day before. The jenny was presented in lateral recumbency. General clinical examination revealed that the Jenny was dull, depressed with vulvaedema and elevated rectal temperature (104 °F). Haematological parameters revealed leucocytosis with neutrophilia. Vaginal examination revealed that fully dilated birth passage engaged with dead

fetus. Further examination of the fetus revealed that the fetus is in oblique ventro vertical presentation (P1), dorso-sacral position (P2) and extended both fore limbs with extension of both the hind limbs beneath the body of fetus and head of the fetus touches the breast region of the fetus (P3). Hence the case was diagnosed as dystocia due to dog sitting posture with nape presentation.

Epidural anaesthesia was given using 3 ml of 2% lignocaine hydrochloride in sacro-coccygeal space. Prior to mutational operation the birth canal was thoroughly lubricated using liquid paraffin, both the forelimbs were secured using obstetrical snare. Repulsion of both the hind limbs and forelimbs were done one by one into the pelvic cavity. Mutational operations were performed to correct the nape presentation; briefly, the fetus was repelled into the gravid uterus to correct the neck deviation after creating the pelvic room space the postural defect of head and neck was corrected. Then William's long obstetrical hook placed on the inner canthus of the left eye of the fetus and traction applied to bring the fetal head out of vulva. Both the fore limbs were taken outside the vulva by gentle traction. Finally, a dead male foal along with fetal membranes delivered per-vaginally. As a part of post obstetrical management therapy the jenny was treated with parenteral fluids, antibiotics, anti-inflammatory and anti-histamines as follows for five days. Inj. Ringer's Lactate 1 lit i.v., Inj. Dextrose Normal Saline 1 lit i.v., Inj.

*Corresponding author: jayatamil9777@gmail.com



Fig. 1. Dead male foal with fetal membrane after delivery

Ceftriaxone 20 mg/kg i.v., Inj. Oxytocin 30 I.U. i.m., Inj. Flunixin meglumine 1.1 mg/kg i.v., Inj. Tetanus toxoid-30 LF units i.m., Inj. Chlorpheniramine maleate 0.5 mg/kg i.m.. The animal recovered uneventfully without much complications.

Dystocia in equids is often considered to be the life-threatening disorders for both the mare and foal. There is a strong association exist between the duration of stage II labour and foal survival; hence, the necessary obstetrical interventions need to be taken to safeguard the life of the fetus and to the prevent serious injuries to the dam. So, action must be taken as soon as a problem is detected (Vaala, 2006). If a mare persists in 2nd stage of labour

beyond 20 minutes without fetal movement dystocia must be suspected (Frazer, 2009). Further, caesarean section in equines is indicated for irreducible fetus, severe maldisposition and protracted dystocia (Dutt *et al.*, 2020). In our case, nape presentation and dog sitting posture might be due to improper orientation of fetus prior to foaling or might be due to improper extension and flexion of fetal extremities.

REFERENCES

- Arthur, G.H., Noakes, D.E. and Pearson, H. (2001). Veterinary reproduction and obstetrics. (6th Edn.), ELBS, Bailliere Tindall, London, UK. pp. 175.
- Baldwin, J.L., Cooper, W.L. and Vanderwall, D.K. (1991). Dystocia due to anterior presentation with unilateral or bilateral hip flexion posture (dog sitting presentation) in the mare. Incidence management and outcomes. In: *Proc 37th Annual Meet Am. Assoc. Eq. Pract.*, pp. 229-40.
- Card, C. (2002). Dystocia in the mare. *Large Anim. Vet. Rounds*. 2: 1-6.
- Dutt, R., Niwas, R., Kumar, S., Dinesh, Saini, D., Dangi, A. and Poonia, A. (2020). Surgical management of dystocia due to foetal deformity in a mare. *Vet. Res. Intern.* 8(1): 41-43.
- Frazer, G.S. (2009). Dystocia management on the farm. In: *Proc. 15th SIVE Congress Annual Meet, Italian Assoc. Equine Vet.*, pp. 57-59.
- Noakes, D.E. (2001). Dystocia and other disorder associated with parturition. In; Noakes, D.E., Parkinson, T.J., England, G.C.W., Edn., Arthur's Veterinary Reproduction and Obstetrics, (8th Edn.), Elsevier Ltd., pp. 341-55.
- Vaala, W.E. (2006). Perinatology. In: *The Equine Manual*, Higgins, A.J., Snyder, J.R., Eds.; W.B. (2nd Edn.), Saunders: Philadelphia, PA, USA, pp. 803-804.

THE HARYANA VETERINARIAN

Editors/Editorial Board Members are highly thankful to all the distinguished referees who helped us in the evaluation of articles. We request them to continue to extend their co-operation and be prompt in future to give their valuable comments on the articles for timely publication of the journal.