

MANAGEMENT AND FEEDING PRACTICES DURING TRANSITION PERIOD OF BUFFALO IN THE VINDHYA REGION OF MADHYA PRADESH

SUMAT KUMAR SHAKYA, NAWAL SINGH RAWAT, ANJANI KUMAR MISHRA, SUMAN SANT¹,
KUMAR GOVIL^{2*}, RAJEEV RANJAN³ and KARISHMA KAKOTIYA

Department of Livestock Production and Management,

¹Department of Veterinary and Animal Husbandry Extension Education,

²Department of Animal Nutrition, ³Department of Veterinary Pharmacology and Toxicology,
College of Veterinary Science and Animal Husbandry, Rewa-486001, M.P., India

Received: 22.08.2023; Accepted: 15.10.2023

ABSTRACT

A study was conducted to collect the information on existing management practices and feeding management of buffaloes of the Vindhya region in Madhya Pradesh. Data were collected from randomly selected 240 dairy buffalo owners through personal interview with the help of pre-tested structured schedule. None of the farmer incorporated common salt and mineral mixture in animal ration; however maximum respondent's occasionally added common salt (83.75%) and mineral mixture (34.17%) in the ration during transition period. Farmers in the study area following conventional housing system and only 22.5% and 67.50% buffalo owners practiced to separate their pregnant animals from the herd and propercutting of naval cord after parturition, respectively. The disease incidence occurs in majority of animals during transition period and having vaginal discharge after 14 days. The majority of respondents practiced additional concentrate feeding before and after calving. It can be concluded from the study that in order to increase the milk production in rural conditions there is need to educate the buffalo farmers about importance of salt and mineral mixture, feeding of scientifically balanced ration during transition period with regular vaccination, proper sanitation and segregation of sick and pregnant buffaloes from healthy ones.

Keywords: Buffalo, Feeding, Health, Nutritional status, Transition period

How to cite: Shakya, S.K., Rawat, N.S., Mishra, A.K., Sant, S., Govil, K., Ranjan, R. and Kakotiya, K. (2024). Management and feeding practices during transition period of buffalo in the Vindhya region of Madhya Pradesh. *Haryana Vet.* 63(1): 44-51.

In terms of revenue, employment, and foreign exchange gains, the livestock industry plays a significant role in India's economy. The total population number of bovines (cattle and buffalo) and buffalo population are 302.34 million and 109.85 million according to the 20th Livestock Census. India is largest milk producers in the world. Dairy sector in the India has shown remarkable development in the past decade. The milk production during 2021-22 is 221.06 million tones and growing at an annual growth of 5.29%. The per capita availability of milk is increased to 444 grams/day in 2021-22 (DAHD, 2022-23).

Buffalo can be considered a backbone for the development of dairy industry in India, contributing 56% of the total milk, while, they constitute only 34.6% of total bovine population (FAO, 2012). River-buffaloes are found mainly in South-Asian countries like India and Pakistan, reared mainly for milk and meat production. Buffalo meat is lean and contains less saturated fat compared to beef and pork and having good nutritive value. (Murthy and Davadason, 2003). About 63% of the world's buffalo milk and 95% of buffalo milk in Asia is contributed by Indian buffaloes (Boopathi *et al.*, 2021).

Buffalo milk is healthy as it is richer in saturated fatty acids; it contains higher total solids useful for preparation

of cheese, butter, sweets and ice creams. Buffalo have been used for draft and termed as the 'living tractor'. In addition buffaloes are also widely used to plow or level land, plant crops, puddle rice fields, cultivate crops, pump water, haul carts and shallow draft boats, carry people, thresh grain, press sugar cane, haul logs, and much more for farmers particularly marginal farmers.

The transition period is the period from 3 weeks pre-calving to 3 weeks post-calving (Grummer, 1995). The term transition defines the important physiological, metabolic and nutritional changes occurring in this time duration. This phase constitutes a turning point in the productive cycle of the cow going to next lactation. Most farmers often ignore this crucial period in productive lifecycle of the animal leading to decreased milk production in the coming lactation and higher incidence of metabolic disorders. Thakur *et al.* (2016) studied the nutrient status of buffaloes of Indore district of Malwa region of Madhya Pradesh and found shortage of DCP, P, Zn, Cu, vitamin A and vitamin E by 11%, 21%, 65%, 2%, 45% and 80%, respectively in advanced pregnant buffaloes.

The nitrogen utilization and growth in Swamp buffalo is more efficient than Malaysian cattle (Devendra, 1985). This better nitrogen utilization and growth is particularly noticeable when the feed supply is of low quantity and/or quality (Wanapat *et al.*, 2003). Praveen and

*Corresponding author: kumargovil22@gmail.com

Dhaarani, (2018) found that failure in proper management during transition phase leads to lowered milk production, immune suppression, poor reproductive performance. Therefore, keeping in view the aforesaid, it is necessary to acquire the data on management practices and feeding management of the buffaloes during transition period in Vindhya region of Madhya Pradesh.

MATERIAL AND METHODS

The present study was conducted in Vindhya region of Madhya Pradesh state. As per Livestock Census 2019, total population of Buffalo in Vindhya region of Madhya Pradesh is 10.3 Million. Four districts (Rewa, Satana, Sidhi and Singrauli) were selected from Vindhya region of Madhya Pradesh. Three tehsils and two villages from each district were randomly selected from identified district. Afterwards, preliminary survey was conducted to know the numbers of farmers practicing the buffalo rearing.

Table 1. Selected Tehsils and Villages of Rewa district

District	Tehshil	Village
Rewa	Teonther	Maghigawan
		Barha
	Jawa	Banigawan
	Hanumana	Janakhaikalan
Satna	Nagod	Khatkhari
		Gauri
		Semarwara
	Maihar	Bamhaur
	Raghuraj Nagar	Podi
Sidhi	Sihawal	Bharauli
		Rampur
	Gopdbanas	Guluwa
		Bithauli
Singrauli	Chitrangi	Bichhri
		Upani
	Deosar	Bhatha
		Tala
Waidhan	Saraiha	Pondi (Bagdari)
		Kusahi
	Jhara	
	Jiawan	
	Harrahawa	
	Tiyara	

From selected villages randomly 10 buffalo owners as respondents were chosen randomly from each selected villages, thus, data from total 240 dairy farmers were collected for management and feeding practices adopted by the farmers of the selected area. The questionnaire regarding management practices and feeding management was developed and pilot tested prior to the implementation. The farmer's houses were regularly visited to gather the data mostly during the morning hours. The data obtained

were analyzed to determine the frequency and percentage.

- a. Study on management practices of buffaloes during transition period-

Table 2. Management practices during transition period

S.No.	Parameters	Measurement
1.	Segregation of pregnant buffalo from the herd (one month before expected date of calving)	Through interview schedule
2.	Calving assistance required for buffaloes	
3.	Naval cord disinfection in buffalo calves	
4.	The disease incidence in buffaloes	
5.	Vaginal discharge after 14 days of calving in buffaloes	
6.	Sanitary practices inside buffalo shed	

- b. Feeding management studied during transition period are as below-

Table 3. Feeding management during transition period

S.No.	Parameters	Measurement
1	Feeding of concentrate to advance pregnant heifer	Through interview schedule
2	Special feeding or extra amount of concentrate to the buffaloes after calving	
3	Concentrate feeding based on feeding standard	
4	Challenge feeding for buffaloes	

RESULT AND DISCUSSION

- a. **Management practices during transition period**

The information regarding management practices during transition period opted by the buffalo owners in Rewa, Satna, Sidhi and Singrauli districts are discussed as below-

1. **Segregation of pregnant buffaloes from the herd**

All the farmers rearing the animal under conventional housing system. On the basis of data obtained it was reported that 77.50% of the respondents did not separated and 22.50% respondents separated pregnant buffalo one month before the expected date of calving from the herd (Table 4). All of the respondents housed sick and healthy buffaloes together may be due to lack of knowledge about the isolation and segregation process to control the spread of diseases in the herd or due to less availability of space.

The district wise results of present investigation was also observed that majority 96.67%, 61.67%, 95.00% and 56.67% of the respondents did not separate of pregnant buffalo one month before the expected date of calving from the herd, whereas only 3.33%, 38.33%, 5.00% and 43.33% of the respondents practices the separation of pregnant buffalo from the herd in Rewa, Satna, Sidhi and Singrauli district, respectively.

2. Calving assistance required for buffaloes

On the basis of data obtained it is found that overall 91.25% of respondents did not need any assistance during calving, whereas only 8.75% of the respondents needed assistance during calving of buffalo as shown in Table 4.

The district wise results of present investigation in case of requirement of assistance during calving, the majority 93.33%, 98.33%, 81.67% and 91.67% of respondents did not need any assistance during calving, whereas only 6.67%, 1.67%, 18.33% and 8.33% of the respondents needed assistance during calving of buffaloes in Rewa, Satna, Sidhi and Singrauli district, respectively. Similarly, Kumar and Mishra (2011), Bais and Singh (2013) and Sabapara *et al.* (2015) also reported assistance during calving amongst majority of farmers at the time of parturition. However, contrary to the present findings by Pata *et al.* (2019) reported majority of respondents attended calving and took care of the calves after parturition in the Junagadh and Porbandar districts of Gujarat.

3. Naval cord disinfection in buffalo calves

The findings presented in Table 4 revealed that overall 67.50% of the respondents followed the navel disinfection of calf after birth, while 32.50% of respondents not practicing the proper ligation, cutting and disinfection of the navel cord leaving it to fall off naturally. The higher percentage of dairy farmers following these practices of navel cord disinfection was probably due to awareness. The district wise results of present investigation revealed that majority 100.00%, 53.33%, 66.67% and 50.00%, of the respondents practiced navel cord disinfection in calf after parturition, whereas 0.00%, 46.67%, 33.33% and 50.00% of the respondents did nothing for navel cord disinfection in Rewa, Satna, Sidhi and Singrauli district, respectively.

For proper navel cord disinfection, it should be cut two inches skin, the contents is squeezed out, dipped in iodine tincture and tied using clean thread to prevent infection. In the present study, majority of the farmers were aware about the navel cord disinfection and care of new born, it helped to reduce the calf mortality. These findings are well comparable with finding of Bilal *et al.* (2019), who reported majority of the farmers practicing navel cord cutting and disinfection. There still is room for improvement in this regard, who are not practicing navel cord care are exposing their newborn calves to diseases. Contrary to our findings, Pawar *et al.* (2006), Sabapara *et al.* (2010) and Kour *et al.* (2019), they revealed that majority of the respondents did not practice navel cord disinfection and it was left to fall off itself naturally.

4. The disease incidence in buffaloes

The findings presented in Table 4 revealed that the in Vindhya region overall 90.84% of the respondents told that the disease incidence occurs during transition period, whereas only 9.16% of the respondents told that buffalo escaped from the diseased incidence during transition period. Among disease incidence in buffaloes in the Vindhya region, majority of buffaloes were suffering from mastitis, retained fetal membranes, metritis, hypocalcaemia, dystocia, prolapse 57.79%, 30.28%, 6.43%, 2.75%, 2.29% and 0.46%, respectively were reported. The district wise results of present investigation reported that the disease incidence occurs during transition period in majority 78.33%, 100.00%, 85.00% and 100.00% of buffaloes, whereas only 21.67%, 0.00%, 15.00% and 0.00% of buffaloes escaped from the diseased incidence in Rewa, Satna, Sidhi and Singrauli districts, respectively.

Among disease incidence in buffaloes, majority of buffaloes were suffering from mastitis, retained fetal membranes, prolapse and dystocia in different proportion in above mention district. The high incidence of disease occurrence might be attributed to poor managerial practices followed by the farmers. The results showed that farmers were unaware about most of the infectious disease occurring during pregnancy and calving. Similarly, Thakur *et al.* (2017) and Pata *et al.* (2019) reported that metabolic disorders are the major problem in buffalo farming, followed by mastitis and reproductive disorders. Scientific management of dairy buffaloes also helped to prevent the disease incidence.

5. Vaginal discharge after 14 days of calving in buffaloes

The findings presented in Table 4 revealed that 100.00% of the buffaloes had vaginal discharge after 14 days in Vindhya region. The overall incidence of white and clear vaginal discharge after 14 days among the buffaloes were found to be 75% and 25%, respectively. The district wise results of present investigation revealed that majority about 100.00%, 100.00%, 100.00% and 100.00% of the buffaloes had vaginal discharge after 14 days in Rewa, Satna, Sidhi and Singrauli district, respectively. Among these buffaloes, 66.67%, 75.00%, 81.67% and 76.67% showed white discharge, while only 33.33%, 25.00%, 18.33% and 23.34% showed clear vaginal discharge after 14 days in Rewa, Satna, Sidhi and Singrauli district, respectively.

According to Sheldon *et al.* (2006) approximately 80 to 100% of cattles have bacteria in their uterine lumen within the first 2 week after calving. Although, the immune system of the body gradually removes the microbes, still up to 40% of cows have bacterial contamination in their

Table 4. Management practices followed by buffalo owners during transition period**1. Segregation of pregnant buffaloes from the herd**

Category	Rewa		Satna		Sidhi		Singrauli		Overall	
	N	%	N	%	N	N	%	N	%	N
Yes	2	3.33	23	38.33	3	5.00	26	43.33	54	22.50
No	58	96.67	37	61.67	57	95.00	34	56.67	186	77.50
2. Calving assistance required for buffaloes										
Yes	4	6.67	1	1.67	11	18.33	5	8.33	21	8.75
No	56	93.33	59	98.33	49	81.67	55	91.67	219	91.25
3. Naval cord disinfection in buffalo calves										
Yes	60	100.0	32	53.33	40	66.67	30	50.00	162	67.50
No	0	0.00	28	46.67	20	33.33	30	50.00	78	32.50
4. The disease incidence in buffaloes										
Yes	47	78.33	60	100.00	51	85	60	100.00	218	90.84
No	13	21.67	0	0.00	9	15	0	0.00	22	9.16
Lameness	0	0.00	0	0.000	0	0	0	0.00	0	0.00
Metritis	0	0.00	4	6.67	0	0	10	16.67	14	6.43
Mastitis	5	10.64	55	91.66	21	35	45	75.00	126	57.79
Hypocalcaemia	0	0.00	1	1.67	0	0.00	5	8.33	6	2.75
ROP	36	76.6	0	0.00	30	50	0	0.00	66	30.28
Prolapse	1	2.12	0	0.00	0	0.00	0	0.00	1	0.46
Dystocia	5	10.64	0	0.00	0	0.00	0	0.00	5	2.29
5. Vaginal discharge after 14 days of calving in buffaloes										
Yes	60	100.0	60	100.0	60	100.0	60	100.0	240	100
No	0	0.00	0	0.00	0	0.00	0	0.00	00	0.00
White	40	66.67	45	75.00	49	81.67	46	76.67	180	75.00
Thick	0	0.00	0	0.00	0	0.00	0	0.00	00	0.00
Sticky	0	0.00	0	0.00	0	0.00	0	0.00	00	0.00
Red	0	0.00	0	0.00	0	0.00	0	0.00	00	0.00
Clear	20	33.33	15	25.00	11	18.33	14	23.33	60	25.00
6. Sanitary practices inside buffalo shed										
Yes	24	40.00	45	75.00	42	70.00	37	61.67	148	61.67
No	36	60.00	15	25.00	18	30.00	19	38.33	92	38.33
Bleaching powder	7	29.17	5	11.10	28	66.67	4	10.81	44	29.73
Phenol	14	58.33	20	44.45	12	28.57	14	37.83	60	40.54
Detergent	3	12.50	20	44.45	2	4.76	19	51.36	44	29.72

(N and % indicate the number and percentage of respondents, respectively)

uterus 3 week after parturition (Sheldon *et al.*, 2008). The bacteria in the uterus and the acute-phase protein response causing clinical endometritis, thus, vaginal discharge (Williams *et al.*, 2005).

6. Sanitary practices inside buffaloes shed

Results can be observed from the Table 4 showed that overall 61.67% of the respondents practiced different measure to sanitize the buffalo shed whereas only 38.33% of the respondents were not followed any measure sanitize the buffalo shed in study area. The present study revealed

that about 40.00%, 75.00%, 70.00% and 61.67% of farmers practiced different measure to sanitize the shed whereas, 60.00%, 25.00%, 30.00% and 38.33% of farmers did not practiced any sanitization process in Rewa, Satna, Sidhi and Singrauli district, respectively.

The district wise results of present investigation showed that among the different sanitization practice, about 29.17%, 11.10%, 66.67% and 10.81% of farmers used Bleaching powder, whereas 58.33%, 44.45%, 28.57% and 37.83% of farmers used Phenol and 12.50%,

Table 5. Feeding management studied during transition period**1. Feeding of concentrate to advance pregnant heifer**

Category	Rewa		Satna		Sidhi		Singrauli		Overall	
	N	%	N	%	N	N	%	N	%	N
Without special feeding	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00
For last 15 days	23	38.33	10	16.67	10	16.67	15	25.00	58	24.17
For last one months	34	56.67	50	83.33	40	66.66	45	75.00	169	70.42
For last two months	3	5.00	0	0.00	10	16.67	0	0.00	13	5.41
2. Special feeding to the buffaloes after calving										
Special feeding	10	16.67	56	93.33	10	16.67	40	66.67	116	48.34
Without special feeding	50	83.33	4	6.67	50	83.33	20	33.33	124	51.66
3. Concentrate feeding based on feeding standard										
Feeding standard	28	46.67	0	0.00	10	16.67	0	0.00	38	15.84
Without feeding standard	32	53.33	60	100.0	50	83.33	60	100.0	202	84.16
4. Challenge feeding for buffaloes										
Challenge feeding	38	63.33	51	85.00	33	55.00	48	80.00	170	70.83
Without challenge feeding	22	36.67	9	15.00	27	45.00	12	20.00	49	29.27

(N and % indicate the number and percentage of respondents, respectively)

44.45%, 4.76% and 51.36% of farmers used Detergent to sanitize the shed in Rewa, Satna, Sidhi and Singrauli district, respectively. This might be due to that the farmers aware about sanitary and hygienic condition in buffalo shed. Regarding the sanitary condition of shed, the present findings are comparable to Aulakh and Rajbir (2012) and Viswkarma *et al.* (2018) who reported that majority of the farmers used sanitizer and disinfectant for cleaning of their shed. However, Khadda *et al.* (2017) and Vranda *et al.* (2017), reported that majority of farmers, did not give more attention towards sanitary condition of shed and not using sanitizer and disinfectant for cleaning of their shed.

b. Feeding management during transition period

The information regarding nutritional status during transition period opted by the buffalo owners in Rewa, Satna, Sidhi and Singrauli districts. The different nutritional managemental practices during transition period related to buffalo were discussed with following heads-

1. Feeding of concentrate to advance pregnant heifer

The results of the Table 5 indicated that overall highest 70.42% of the buffalo keepers were feeding concentrate to heifers for last one month during advance pregnancy, While 24.17% of the respondents provide concentrate feed for last 15 days and only 5.41% of the respondents provide concentrate feed for last two month to

advance pregnant heifer during transition period. The district wise results of present investigation reported that majority 56.67%, 83.33%, 66.66% and 75.00% of respondents provide concentrate feed for last one month during advance pregnancy of heifer, while 38.33%, 16.67%, 16.67% and 25.00% of the respondents provide concentrate feed for last 15 days and only 5.00%, 0.00%, 16.67% and 0.00% of the respondents provide concentrate feed for last two month in Rewa, Satna, Sidhi and Singrauli district, respectively.

Scientific feeding during transition phase is very important as maximum development of fetus occurs during last trimester of pregnancy. The supplementation of mineral mixture to the pregnant buffaloes improves productive performance of the buffaloes. The digestive system of high yielders has better concentrate digestion resulting in body weight gain and improvement of body condition of buffaloes.

Similarly, Chowdhry *et al.* (2006), Rathore and Kachwaha (2009), Sabapara *et al.* (2016), Kumar and Mishra (2011), Kumar *et al.* (2011), Manohar *et al.* (2014) and Kumar *et al.* (2019) reported that majority of the buffaloes farmers feeding concentrate to the advanced pregnant heifers. Contrary, Madke *et al.* (2006), Kishore *et al.* (2013) and Rangamma *et al.* (2013) reported few farmers feeding concentrate to their advanced pregnant buffaloes. Sabapara *et al.*, 2016 and Kumar *et al.* (2019)

also found that majority percent of the respondents feeding concentrates to their advanced pregnant heifers during last 2 months of pregnancy.

2. Special feeding to the buffaloes after calving

The results of the Table 5 indicated that overall about 48.34% of respondents practiced special feeding after calving, while 51.66% of respondents did not practiced any type of special feeding after calving. The district wise results of present investigation indicated that 16.67%, 93.33%, 16.67% and 66.67% of respondents practiced special feeding after calving whereas, without special feeding after calving was 83.33%, 6.67%, 83.33% and 33.33% in Rewa, Satna, Sidhi and Singrauli districts, respectively.

Most respondents had adequate knowledge about feeding care after calving. These findings are in agreement with Divekar and Saiyed (2008), Sabapara *et al.* (2010), Kishore *et al.* (2013) and Kumar *et al.* (2019), who reported the majority of the respondents, followed special feeding after calving.

3. Concentrate feeding based on feeding standard

The result of present study in table 5, indicated that overall 84.16% of respondent did not followed the guideline of feeding standard, while only 15.84% of respondent nourished their buffalo as per the guideline of feeding slandered and giving concentrate feed based on quantity of milk produced during transition period. The district wise results of present investigation indicated that majority i.e. 53.33%, 100.00%, 83.33% and 100.00% of respondent did not followed the guideline of feeding slandered, while 46.67%, 0.00%, 16.67% and 0.00% of respondent nourished their buffaloes as per the guideline of feeding slandered and giving concentrate feed based on quantity of milk produced in Rewa, Satna, Sidhi and Singrauli, districts, respectively.

Similarly, Dar *et al.* (2017) found that majority of the respondents (97.75%) not feeding animals according to productivity level. Contrary, Malik *et al.* (2005), Divekar and Saiyed (2008), Sheikh *et al.* (2011), Sabapara *et al.* (2016) and Kumar *et al.* (2019), reported that majority of the respondents (87.50%) fed concentrates to their animals as per milk production, and few percent of the respondents fed concentrates to their animals on a no criteria basis. Manohar *et al.* (2014) reported that majority of respondent feeding 1 kg concentrate per 2 litres of milk per day to the lactating buffaloes.

4. Challenge feeding for buffaloes

The results of challenge feeding practices studied for buffaloes were presented in Table 5, showed that

overall about 70.83% of respondents allowed challenge feeding, while 29.27% of the respondents did not allowed challenge feeding to their buffalo.

The district wise results of present investigation showed that majority about 63.33%, 85.00%, 55.00% and 80.00% of respondents allowed challenge feeding, while 36.67%, 15.00%, 45.00% and 20.00% of the respondents did not allowed challenge feeding to their buffaloes in Rewa, Satna, Sidhi and Singrauli district, respectively. These results were in accordance with Patil (1996) and Bourde (1992).

CONCLUSION

On the basis of data obtained from the present study on buffalo farmers of Vindhya region about management and feeding management during transition period of the animal, it is concluded that majority of the farmers did not separate of pregnant buffalo from the herd, did not need any assistance during calving, face the navel disinfection of calf after birth, disease incidence occurs during transition period, and almost every buffalo (100.00%) had vaginal discharge after 14 days, and farmers were practicing different measure to sanitize the buffalo shed. In addition, buffalo keepers were feeding concentrate to heifers for last one month during advance pregnancy but did not followed the guideline of feeding standard during transition period, while, lesser farmers practiced scientific feeding to the animal after calving and allowed challenge feeding during transition period. Thus in order to increase the milk production in rural conditions there is need to educate the buffaloes farmers about importance of salt and mineral mixture in buffaloes ration, feeding of scientifically balanced ration in transition period.

REFERENCES

- Aulakh, G.S and Rajbir, S.(2012). Adoption of recommended management practices by the buffalo owners. *Indian J. Dairy Sci.* **65(5)**: 431-434.
- Bais, B. and Singh, D.M. (2013). Existing buffalo calf rearing management practices in Jaipur district of Rajasthan in India. *J. Vet. Sci. Technol.* **4(4)**: 80.
- Bilal, M., Xu, Y. and Rao, M.I. (2019). Assessment of calf health care and management practices under small holder production system in Punjab. *Pakistan Buffalo Bull.* **38(1)**: 57-66.
- Boopathi, V., Prasad, S., Kumaresan, A., Manimaran, A. and Prakash, M.A. (2021). Environmental factors affecting reproductive performance of Murrah buffaloes. *Biol. Rhythm Res.* **52(6)**: 1-7.
- Bourde, S.V. (1992). An impact of dairy co-operative society on socioeconomic development of member farmers. M.Sc. (Agri.) thesis, M.P.K.V. Rahuri.
- Chowdhary, N.R., Patel, J.B and Bhakat, M. (2006). An overview of feeding, breeding and housing practices of dairy animals under milk co-operative system in Banaskantha district of North Gujarat region. *Dairy Planner.* **5(12)**: 8-10.

- Chowdhary, N.R., Patel, J.B. and Bhakat, M. (2006). An overview of feeding, breeding and housing management practices of dairy animals in the tribal area of South Gujarat. *Indian J. Anim. Sci.* **80(10)**: 1022-1027.
- DAHD (2023). Dairy, Animal Husbandry Department, Government of India, available from. <http://dahd.nic.in/about-us/divisions/cattle-and-dairy-development>.
- Dar, P.A., Azmat Alam Khan, A.A. Shah, A.A., Qadri, I.A. and H.M. Khan (2017). Studies on breeding and feeding practices of buffalo in vogue among buffalo farmers in temperate himalayan region of Kashmir Valley. *Indian J. Hill Farming. (Special Issue)*. pp. 83-87.
- Devendra, C. (1985). Comparative nitrogen utilization in Malaysia swamp buffaloes and Kedah-Kelanton cattle. In: Proc. The 3rd AAAP Animal Science Congress, Seoul, Korea, **2**: 873-875.
- Divekar, B.S. and Saiyed, L.H. (2008). Feeding practices followed by professional cattle owners of Anand district. *Indian J. Field Veterinarians.* **3(4)**: 31-34.
- FAO (2012). <http://faostat.fao.org/site/569/default.aspx#ancor>.
- Grummer, R.R. (1995). Impact of changes in organic nutrient metabolism on feeding the transition dairy cow. *J. Anim. Sci.* **73(9)**: 2820-2833.
- Khadda, B.S., Lata, K., Singh, B. and Kumar, R. (2017). Study of buffalo husbandry practices in rural area of central Gujarat in India. *Buffalo Bulletin.* **36(1)**: 75-87.
- Kishore, K., M. Mahender, and C. Harikrishna (2013). A study on buffalo management practices in Khammam district of Andhra Pradesh. *Buffalo Bulletin.* **32(2)**: 97-119.
- Kour, S., Khan, A., Brahma, B., Jeelani, R., Khursheed, I. and Dua, S. (2019). A study on buffalo management practices followed by Gujjar tribe of Jammu. *J. Pharmacogn. Phytochem.* **8(3)**: 218-221.
- Kumar, J., Singh, R., Somnath, R., Dayal, H. and Singh, S. (2019). Studies on feeding and breeding practices of dairy animal in western Uttar Pradesh. *J. Pharmacogn. Phytochem.* **3**: 29-36.
- Kumar, M. (2015). Buffalo healthcare management practices followed by the farmers of Ferozepur district of Punjab, India. *Indian J. Anim. Res.* **49(3)**: 413-415.
- Kumar, S. and Mishra, B.K. (2011). Existing calf rearing and milking management practices followed by dairy farmers in Uttarakhand. *J. Hill Agric.* **2(1)**: 78-84.
- Kumar, S. and Mishra, B.K. (2011). Existing feeding and housing management practices followed by dairy producers in Tehri Garhwal district of Uttarakhand. *Indian J. Anim. Prod. Mgt.* **27(3-4)**: 159-162.
- Kumar, S., Mishra, B.K., Yadav, J.S and Kumar, A. (2011). Existing breeding and health management practices followed by dairy farmers in mid hills of Uttarakhand. *Indian J. Anim. Prod. Mgt.* **27(1-2)**: 34-37.
- Khadda, B.S., Lata, K., Singh, B. and Kumar, R. (2017). Study of buffalo husbandry practices in rural area of central Gujarat in India. *Buffalo Bulletin.* **36(1)**: 75-87.
- Madke, P.K., Murkute, J.S., Upadhye, S.V. and Vedpathak, C.P. (2006). Adoption of scientific feeding practices by dairy farmers. *Indian J. Anim. Res.* **40(2)**: 155-157.
- Malik, B.S., Meena, B.S. and Rao, S.V.N. (2005). Study of existing dairy farming practices in Uttar Pradesh. *J. Dairy Foods Home Sci.* **24(2)**: 91-95.
- Manohar, D.S., Bais, B., Goswami, S.C., Jhirwal, A.K and Choudhary, D. (2014). Study on breeding management practices of buffaloes in relationship with selected traits of respondents in Jaipur district of Rajasthan (India). *Indian J. Field Vety.* **9(3)**: 82-83.
- Manohar, D.S., Goswami, S.C. and Bais, B. (2014). Study on feeding management practices of buffaloes in relationship with selected traits of respondents in Jaipur district of Rajasthan, India. *Indian J. Anim. Res.* **48(2)**: 150-154.
- Meena, H.R., Ram, H., Sahoo, A. and Rasool, T.J. (2008). Livestock husbandry scenario at high altitude Kumaon Himalaya. *Indian J. Anim. Sci.* **78**: 882-886.
- Murthy, T.R.K. and Devadason, I.P. (2003). Buffalo meat and meat products-An overview. In Proceedings of the 4th Asian Buffalo Congress on Buffalo for Food, Security and Employment: Asian Buffalo Association, New Delhi, India, pp. 193-199.
- Pata, B.A., Odedra, H.H., Savsani, A.R., Ahlawat, T.K. and Odedara, A.B. (2019). Breeding, calf and health management practices opted by buffalo owners in Junagadh and Porbandar districts of Gujarat: A comparative study. *Intl. J. Current Microbiol. Applied Sci.* **8(3)**: 2426-2435.
- Patil, S.S., Mali, S., Land Kalke, S.D. (1996). Preliminary studies on feeding and management practices followed by buffalo owners from Kolhapur district of Maharashtra. National Symposium on Buffalo Research for higher productivity, held at CCS Haryana Agriculture University, Hissar Session- IV. p. 54.
- Pawar, B.K., Nalawade, T.H. and Jagtap, D.Z. (2006). Adoption of bovine feeding practices and constraints faced by tribal farmers of Pune district. *J. Maharashtra Agriculture University.* **31(3)**: 329-330.
- Praveen, S and Dhaarani, C. (2018). Management approach of the cow during transition period. *Int. J. Sci. Environ.* **7(3)**: 950-954.
- Rangamma, B., Jadeswararao, S., Prasad, R.M.V. and Ragavarao, E. (2013). Managemental practices adopted by buffalo milk producers in Krishna district of Andhra Pradesh. *Indian J. Anim. Prod. Mgt.* **29(1-2)**: 61-68.
- Rathore, R.S. and Kachwaha, R.N. (2009). Studies on existing management practices followed by the buffalo owners in Jhunjhunu district of Rajasthan. *Indian J. Anim. Prod. Mgt.* **25(1-2)**: 8-11.
- Sabapara, G.P., Desai, P.M., Kharadi, V.B. and Singh, R.R. (2010). Breeding and health care management status of dairy animals in the tribal area of South Gujarat. *Indian J. Anim. Sci.* **80(11)**: 1148-1151.
- Sabapara, G.P., Desai, P.M., Kharadi, V.B. and Singh, R.R. (2010a). Housing and feeding management practices of dairy animals in the tribal area of South Gujarat. *Indian J. Anim. Sci.* **80**: 1022-27.
- Sabapara, G.P., Fulsoundar, A. and Kharadi, V.B. (2015). Survey of calf rearing practices followed at rural dairy farms in Surat district. *J. Anim. Res.* **5(2)**: 257-261.
- Sabapara, G.P., Fulsoundar, A.B. and Kharadi, V.B. (2016). Profile of dairy farmers and relationship with adoption of improved dairy husbandry practices in southern Gujarat, India. *Livest. Res. Int.* **4(1)**: 36-40.
- Sheikh, A.S., Bhati, D.S. and Sheikh, W. (2011). Feeding practices followed by professional Kankrej cow owners of Banaskantha district of North Gujarat. *Progress. Agric.* **2(1)**: 67-69.
- Sheldon, I.M., Lewis, G.S., LeBlanc, S. and Gilbert, R.O. (2006).

- Defining postpartum uterine disease in cattle. *Theriogenol.* **65(8)**: 1516-1530.
- Sheldon, I.M., Williams, E.J., Miller, A.N., Nash, D.M. and Herath, S. (2008). Uterine diseases in cattle after parturition. *Vet. J.* **176(1)**: 115-121.
- Singh, R.R., Prajapati, V.S., Kharadi, V.B. and Chaudhary, S.S. (2015). Status of breeding and health care management practices of dairy bovines in the rural and urban areas of south Gujarat of India. *J. Anim. Sci. Adv.* **5(11)**: 1514.
- Snedecor, G.W. and W.G. Cochran (1994). *Statistical Methods* (6th Edn.), Oxford and IBH Publishing Co., New Delhi.
- Tewari, H., Kumar, S., Singh, D.V., Rath, R. and Tyagi, K. (2018). Studies on existing milking and health care practices adopted by dairy farmers in Tarai region of Uttarakhand, India. *Indian J. Anim. Res.* **52(3)**: 454-458.
- Thakur, D., Jain, R.K., Sharma, P. and Yadav, A. (2017). Feeding patterns, nutritional status of available feeds during advanced pregnancy and incidence of reproductive and metabolic disorders in buffaloes of indore district of Madhya Pradesh. *Indian J. Anim. Nutri.* **34**: 50-55.
- Vishwkarma, R., Singh, R., Kushram, P., Singh, S.K. and Sharma, S. (2018). Existing status of buffalo husbandry practices in Jabalpur. *Pharm. Innov. J.* **7(2)**: 08-11.
- Vranda, R., Satyanarayan, K., Jagadeeswary, V., Veeranna, K.C., Rajeshwari, Y.B., Vranda, R.K., Jagadeeswary, V. and Shilpa, J. (2017). Health care management practices by buffalo rearing farmers in Karnataka. *Int. J. Livest. Res.* **7(5)**: 168-174.
- Vranda, R.K., Satyanarayan, V., Jagadeeswary, K.C., Veeranna, Y.B., Rajeshwari, G. and Shree, J.S. (2017). A study on different housing practices of buffaloes in Bidar district of Karnataka. *Int. J. Environ. Sci. Technol.* **6(1)**: 295-302.
- Wanapat, M., Nontaso, N., Yuangklang, C., Wora-Anu, S., Ngarmsang, A., Wachirapakorn, C., and Rowlinson, P. (2003). Comparative study between swamp buffalo and native cattle in feed digestibility and potential transfer of buffalo rumen digesta into cattle. *Asian-Australasian J. Anim. Sci.* **16(4)**: 504-510.
- Williams, E.J., Fischer, D.P., Pfeiffer, D.U., England, G.C., Noakes, D.E., Dobson, H. and Sheldon, I.M. (2005). Clinical evaluation of postpartum vaginal mucus reflects uterine bacterial infection and the immune response in cattle. *Theriogenol.* **63(1)**: 102-117.

CONTRIBUTORS MAY NOTE

- Research/Clinical articles are invited for next issue from the Scientists/Veterinarians engaged in Veterinary Profession.
- Please follow strictly the format of 'The Haryana Veterinarian' for manuscript writing/submission.
- Please pay processing fee of Rs. 1000/- online in the account of Dean, College of Veterinary Sciences, along with each article.
- After revision, please return the revised manuscript and rebuttal at the earliest.
- Please mention your article reference number in all correspondence for a quick response.
- We solicit your co-operation.
- All correspondence should be addressed to 'The Editor', Haryana Veterinarian, Department of Veterinary Parasitology, College of Veterinary Sciences, LUVAS, Hisar-125004.

Editors