

ITS2 TARGETED MOLECULAR CHARACTERIZATION AND PHYLOGENETIC ANALYSIS OF CAPRINE AMPHISTOMES FROM UTTAR PRADESH (NORTH INDIA)

AMIT KUMAR JAISWAL*, DAYA SHANKER, AMIT SINGH¹, PRADEEP KUMAR¹,
VIVEK AGRAWAL² and SHANKER KUMAR SINGH³

Department of Veterinary Parasitology, ³Department of Veterinary Medicine,
COVS&AH, DUVASU, Mathura-281001, Uttar Pradesh, India

¹Department of Veterinary Parasitology, COVS&AH, NDUAT, Kumarganj, Ayodhya-224229 (U.P.), India

²Department of Veterinary Parasitology, COVS&AH, Mhow, NDVSU, Jabalpur-482001 (M.P.), India

Received: 30.08.2022; Accepted: 11.12.2022

ABSTRACT

The present study aimed for molecular characterization and phylogenetic analysis of caprine amphistomes in the northern part of India. A total of 92 rumens from slaughtered goats naturally infected with amphistome species were collected. Based on the morphological characters, three major species of amphistomes viz. *Paramphistomum epiclitum*, *Gastrothylax crumenifer* and *Fischoederius* spp. were identified. Randomly selected amphistomes were pressed in between glass slides and examined microscopically and having similar morphology with identified amphistomes were subjected to molecular characterization by amplification of ITS2 DNA region. Genes of Mathura isolates were aligned with previously published sequences of amphistome species from different parts of globe. Multiple sequence alignment reports and phylogenetic analysis with previously published sequences throughout the globe suggested, the minimum evolutionary divergence between sequences of *P. epiclitum* Mathura isolates (KX657873; KX657875; KX657874) and Indian isolates (KF564870) without showing any difference of goat or buffalo origin. Amplified nucleotide sequence of *Fischoederius* spp. of Mathura isolate showed 86.5% phylogeny similarity with *G. crumenifer* India isolate of cattle. The *G. crumenifer* Mathura isolate showed 96.1% phylogeny similarity with *Calcicophoron microbothrium* of Egypt. The percent identity matrix revealed the *Fischoederius* spp. of Mathura isolate showed 87.2% identity with *F. elongatus* India isolate from *Bos indicus* (JQ688410). The *P. epiclitum* of present study showed 74.0% identity with *P. epiclitum* India isolate from *Bubalus bubalis*. The Percent identity of *G. crumenifer* of present study showed 95.6% with *G. crumenifer* India isolate of *Bos indicus*.

Keywords: Goat amphistomes, ITS-2, Molecular characterization, Phylogenetic analysis

How to cite: Jaiswal, A.K., Shanker, D., Singh, A., Kumar, P., Agrawal, V. and Singh, S.K. (2023). ITS2 targeted molecular characterization and phylogenetic analysis of caprine amphistomes from Uttar Pradesh (North India). *Haryana Vet.* 62(SI-2): 22-25.

Among the various parasite of goats, amphistomosis, caused by digenetic trematode (fluke) of the superfamily Paramphistomoidea is one of the least explored trematodal infectious diseases Yamaguti (1971) listed 62 species of paramphistomes of domestic ruminants from various countries. Paramphistomosis is caused by specific species of the parasite in a particular region (Anuracpreeda *et al.*, 2008) with different pathogenicity (Mehlhorn, 2008). Among them, most prevalent amphistomes in India are *P. epiclitum*, *G. crumenifer*, *Fischoederius elongatus*, *F. cobboldi* and *Gigantocotyle explanatum* with the prevalence ranging 50-70% (Prasad and Varma, 1999). In Indian subcontinent, immature paramphistomosis of domestic ruminant ranks next to fasciolosis and the mortality can reach up to 88% in sheep and goats (Agrawal, 2003). The conventional method for identification of these parasites is only based upon morphology. But it is stated that various genera of family Paramphistomatidae are difficult to identify on a morphological basis alone (Horak, 1971). Second internal transcribed spacer (ITS2) of ribosomal DNA sequence is a highly conserved region due to high degree of an inter-specific and low level of intra-

specific genetic variations among populations from different hosts and geographical localities at generic and species level (Choudhary *et al.*, 2015). PCR-based techniques providing ITS2 sequences have proven to be a reliable tool to identify digenetic species and their phylogenetic relationships (Mohanta *et al.*, 2017). The present study explores the molecular characterization and phylogenetic analysis of caprine amphistome of north India origin.

MATERIALS AND METHODS

Collection, Processing, Preservation and morphological identification of parasites

The 92 rumen of goats (infected with amphistome species) were collected in clean polythene bags from a slaughter house in Mathura, Uttar Pradesh, India, and were transported with ice packs immediately to Department of Parasitology, College of Veterinary Science & Animal Husbandry, Mathura (Uttar Pradesh), India for further processing. The slaughter house at Mathura has a catchment of goats from various parts of northern India. The adult amphistomes are localized in the rumen and generally mono species of amphistome is localized in a specific part of rumen at a time. Amphistomes were

*Corresponding author: drakjaiswal@gmail.com

removed gently from each rumen with the help of forceps and transferred to a beaker containing Phosphate Buffered Saline (PBS) and washed several times with PBS solution to remove ruminal contents. Two cleaned parasites with similar in gross morphological characters from the localized area of rumen were selected. They were gently pressed in between glass slides to confirm the similarity in morphology. Among them one glass slide pressed parasite was stored at -20 °C for DNA isolation and molecular characterization while rest one parasites were processed for morphological identification by staining procedure. The preservation and morphological identification of amphistomes were done as per protocol (Say, 1990).

Molecular characterization of amphistomes

Previously published primers for the partial sequence of ITS2 region of amphistomes were custom synthesized using nucleotide pattern from a highly conserved region (GenBank Accession Number JF834888). Details position of nucleotides, nucleotide sequences, and expected PCR products of the primer is mentioned in Table 1. Genomic DNA was extracted from four amphistomes using phenol-chloroform protocol (Sambrook and Russel, 2006) with minor modifications. The purity and concentration were estimated by Nanodrop (Eppendorf, Biophotometer, Germany). The optical density ratio 260/280 was observed for estimation of purity of double-stranded DNA. A concentration of DNA was measured by taking A260 value. After quantitative and qualitative analysis, the working concentration of isolated DNA was adjusted to 100ng/μl with nuclease-free water (NFW).

The PCR reaction was set up in 25 μl volume containing 12.5 μl PCR Master Mix ThermoFisher Scientific (0.05/μl Taq DNA polymerase in reaction buffer, 4 mM MgCl₂, 0.4 mM dATP, 0.4 mM dCTP, 0.4 mM dGTP, and 0.4 mM dTTP), 1.5 μl of 10 pmol each primer (ITS 2 F/R), 2 μl of the DNA template and total volume was made up to 25μl using NFW. The thermocycling conditions (BioRad) used for amplification of the ITS 2 gene was initial denaturation at 95 °C for 5 minutes, denaturation at 94 °C at 5 minute, annealing at 54 °C for 45 seconds, extension and final extension at 72 °C for 1 and 10 minutes, respectively. The total of 35 cycles were used for denaturation, annealing and extension.

The amplified amplicons were analyzed by agarose gel electrophoresis in 1.5 % agarose gel (Invitrogen). After that PCR products were purified using the Gene Jet PCR purification kit (Thermo Fisher Scientific, US) as per the manufacturer's protocol.

Gene sequencing and Phylogenetic analysis

Purified PCR products of four parasite samples were


sequenced by outsource sequencing (Thermo Fisher Scientific, US). Nucleotide sequencing was done in one side direction using the same set of primers that were used for DNA amplification. The sequences, hence obtained, were submitted into the NCBI, and corresponding accession numbers were obtained. Thereafter, the sequences were analyzed by online tool Mega 6. A multiple sequence alignment was generated with the Clustal W program within Mega 6 software (Tamura *et al.*, 2013). The phylogenetic relationship of amphistomes with other isolates of across the globe was commutated based on nucleotide sequences of the ITS using Mega 6 software. A phylogenetic tree for ITS2 was reconstructed using maximum likelihood. *Haemonchus contortus* and *Fasciola gigantica* was used as out-group species to root the tree. The phylogeny was analysed by Maximum Likelihood (ML) with bootstrap method using 1000 bootstrap replications. Model adopted for this analysis was Kimura 2-parameter with nucleotide substitution model. The rates among sites selected gamma distributed with 5 number of discrete gamma categories. Here was complete deletion of gaps/missing data treatment. The selected ML Heuristic Method was Nearest-Neighbor-Interchange (NNI) with strong Branch Swap Filter (Tamura *et al.*, 2013).

RESULTS AND DISCUSSION

The collected amphistomes from the rumen were identified as *Paramphistomum epiclitum*, *Gastrothylax crumenifer* and *Fischoederius* spp (Agrawal, 2003). Among 92 rumen samples, 88 times (95.6%) *Paramphistomum epiclitum* was identified, while *Gastrothylax crumenifer* and *Fischoederius* spp were identified in 3 (3.2%) and 1 (1.08%) rumen, respectively. The stored amphistomes at -20 °C from these three groups were subjected to DNA isolation and molecular characterization using the partial sequence of the 28S rRNA, internal transcribed spacer 2 (ITS2) for confirmation of morphology and the amplicons of about 400 bp were obtained by thermocycling (Fig. 1). The sequence analysis of these amplicons revealed with *Paramphistomum epiclitum*, *Gastrothylax crumenifer* and *Fischoederius* spp.

A total of 04 representative amplicons (two from *Paramphistomum epiclitum*, one from each *Gastrothylax crumenifer* and *Fischoederius* spp.) were underwent BLAST to confirm these with morphological identification. Then after amplicons were submitted to the National Centre for Biotechnology Information (NCBI) and accession numbers were obtained (KX657873, KX657875, KX639720 and MN535799) Multiple sequence alignment report suggested that there was few nucleotide variations in Mathura isolate with other Indian isolates (JX678261;

Gene	Primer Sequence		Size of amplicon (bp)
ITS 2	5'GAATTAATGTGAAGTC ATACTGC3'	FORWARD	Approx 400 bp
	5'TGATATGCTTAAGTTCA GCGGG3'	REVERSE	



The minimum evolutionary divergence between sequences were reported in *P. epiclutum* Mathura isolates (KX657873; KX657875; KX657874) and Indian isolates (KF564870) without showing any difference of goat or buffalo origin. The phylogenetic analysis revealed that *P. epiclutum* of Mathura isolates related in the same clad without showing any difference of goat or buffalo origin. The phylogenetic analysis and percent identity matrix of amplified nucleotide sequence of *Fischoederius* spp. of Mathura isolate showed 86.5% phylogeny similarity with *G. crumenifer* India isolate of cattle. The *G. crumenifer* Mathura isolate showed 96.1% phylogeny similarity with *Calcicophoron microbothrium* of Egypt. The percent identity matrix revealed the *Fischoederius* spp. of Mathura isolate showed 87.2% identity with *F. elongates* India isolate from *Bos indicus* (JQ688410). The *P. epiclutum* of present study showed 74.0% identity with *P. epiclutum* India isolate from *Bubalus bubalis*. The Percent identity of *G. crumenifer* of present study showed 95.6% with *G. crumenifer* india isolate of *Bos indicus*.

KC283190 | *G. crumenifer* | *Bos indicus* | Ind
 39
 KX639720 | *Fischoederius* spp. | *Capra hircus* | MTR ▲
 JQ688412 | *G. crumenifer* | *Capra hircus* | Ind
 JQ688410 | *F. elongatus* | *Bos indicus* | Ind
 GU133061 | *F. elongatus* | *Capra hircus* | Ind
 23
 AB817059 | *Calicophoron microbathrium* | *Bos taurus* | Egy
 31
 KX274233 | *P. cervi* | *Cervus elaphus* | Croa
 KP341659 | *P. cervi* | *Bos grunniens* | Chi
 88
 MN535799 | *Gastrothylax crumenifer* | *Capra hircus* | MTR ▲
 KF564870 | *P. epiclitum* | *Bubalus bubalis* | Ind
 48
 31
 KX657874 | *Paramphistomum epiclitum* | *Bubalus bubalis* | MTR
 86
 JX678261 | *P. epiclitum* | *Bos indicus* | Ind
 57
 KX657873 | *Paramphistomum epiclitum* | *Capra hircus* | MTR ▲
 KX657875 | *Paramphistomum epiclitum* | *Capra hircus* | MTR ▲
 81
 KJ995525 | *P. leydeni* | *Bos taurus* | Arg
 AB743577 | *Explanatum explanatum* | *Bubalus bubalis* | Jap
 KY889140 | *Explanatum explanatum* | *Bubalus bubalis* | MTR
 AB553695 | *Fasciola gigantica* | *Bubalus bubalis* | Egy
 FN432335 | *Haemonchus gigantica* | *Capra hircus* | Italy

0.1

	Percent identity																			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	
Divergence	1	0.53	0.21	0.03	0.72	0.25	0.05	0.82	0.52	0.28	0.07	0.82	0.12	0.13	0.73	0.85	0.28	0.68	0.72	1
	2	0.17	0.47	0.71	0.03	0.18	0.18	0.42	0.05	0.42	0.72	0.53	0.28	0.18	0.78	0.17	0.18	0.61	0.52	2
	3	0.47	0.17	0.51	0.05	0.18	0.58	0.58	0.05	0.42	0.72	0.53	0.18	0.07	0.82	0.12	0.03	0.82	0.10	3
	4	0.25	0.03	0.51	0.18	0.78	0.68	0.06	0.94	0.94	0.82	0.72	0.18	0.78	0.12	0.55	0.80	0.84	0.81	4
	5	0.82	0.47	0.18	0.05	0.84	0.75	0.72	0.05	0.58	0.18	0.72	0.08	0.82	0.75	0.18	0.61	0.51	0.51	5
	6	0.25	0.47	0.72	0.18	0.08	0.72	0.05	0.82	0.18	0.72	0.08	0.82	0.75	0.18	0.61	0.51	0.51	0.51	6
	7	0.05	0.18	0.47	0.72	0.18	0.08	0.72	0.05	0.82	0.18	0.72	0.08	0.82	0.75	0.18	0.61	0.51	0.51	7
	8	0.12	0.13	0.07	0.82	0.12	0.03	0.82	0.18	0.72	0.08	0.82	0.75	0.18	0.61	0.51	0.51	0.51	0.51	8
	9	0.18	0.68	0.43	0.58	0.28	0.28	0.42	0.75	0.71	0.72	0.21	0.08	0.05	0.81	0.78	0.40	0.67	0.8	9
	10	0.13	0.71	0.72	0.28	0.28	0.28	0.18	0.78	0.61	0.81	0.21	0.23	0.65	0.80	0.78	0.63	0.8	0.8	10
	11	0.78	0.17	0.72	0.28	0.32	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	11
	12	0.07	0.82	0.72	0.28	0.32	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	12
	13	0.82	0.07	0.72	0.28	0.32	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	13
	14	0.07	0.82	0.72	0.28	0.32	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	14
	15	0.12	0.13	0.07	0.82	0.12	0.03	0.82	0.18	0.72	0.08	0.82	0.75	0.18	0.61	0.51	0.51	0.51	0.51	15
	16	0.73	0.18	0.68	0.43	0.58	0.28	0.28	0.42	0.75	0.71	0.72	0.21	0.08	0.05	0.81	0.78	0.40	0.67	16
	17	0.17	0.72	0.72	0.28	0.32	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	17
	18	0.68	0.17	0.72	0.28	0.32	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	0.28	18
	19	0.72	0.61	0.81	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	0.51	19
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	
	1	0.63	0.																	

amphistomes. It is suggested that primer based on ITS2 is a good marker for identification of species of amphistomes after using too BLAST. It is also confirmed that 28S ribosomal DNA region is a useful molecular marker for species identification and Platyhelminthes phylogeny (Kumar *et al.*, 2014). Moreover, ITS2 has been described as a useful marker for species identification of amphistomes (Goswami *et al.*, 2009). The phylogenetic analysis also revealed that the Mathura isolates were closer to the published sequence of different isolates of amphistomes species of Indian origin without showing difference of goat and buffalo origin. The study also indicated the strain of amphistomes particularly *P. epiclitum* was not host species specific. Despite quite an expensiveness of molecular method, it could be helpful for identification of amphistomes with high accuracy as well as to know the phylogenetic relationship with other isolates. In

conclusion, the present study revealed that the amplification of ITS2 region of amphistomes genomic DNA is a reliable tool to know the phylogenetic relationship among *P. epiclutum*, *G. crumenifer* and *Fiscoederius* sp. using tool BLAST The species of *P. epiclutum* of present study showed a close phylogenetic relationship in between and other Indian isolates without showing difference of goat or buffalo origin. The study indicated that the strain of amphistomes particularly *P. epiclutum* was not a host species specific (Sanguankiat *et al.*, 2016 and Jadhav *et al.*, 2018). Although further investigation is required with large number of samples.

REFERENCES

- Agrawal, M.C. (2003). Epidemiology of fluke infections. In: Sood ML (ed) *Helminthology in India* (1st Edn.), International Book Distributors, Dehradun. pp. 511-542.
- Anuracpreeda, P., Wanichanon, C. and Sobhon, P. (2008). Paramphistomumcervi: antigenic profile of adults as recognized by infected cattle sera. *Experimental Parasitology*. **118**: 203-207.
- Choudhary, K., Verma, A.K., Swaroop, S. and Agrawal, N. (2015). A review on the molecular characterization of digenean parasites using molecular markers with special reference to ITS region. *Helminthologia*. **52**: 167-187.
- Goswami, L.M., Prasad, P.K., Tandon, V. and Chatterjee, A. (2009). Molecular characterization of *Gastrodiscoides hominis* (Platyhelminthes: Trematoda: Digenea) inferred from ITS rDNA sequence analysis. *Parasitology Research*. **104**: 1485-1490.
- Horak, I.G. (1971). Paramphistomiasis of domestic ruminants. *Advances in Parasitology*. **9**: 33-72.
- Jadhav, M.M., Kumar, N., Das, B. and Solanki, J.B. (2018). Morphological and molecular characterization of *Paramphistomum epiclutum* of small ruminants. *Acta Parasitologica*. **63**: 586-594.
- Kumar, A., Chaudhary, A., Verma, C. and Singh, H.S. (2014). Molecular Characterization of *Gastrothylax crumenifer* (Platyhelminthes: Gastrothylacidae) from Goats in the Western Part of India by LSU of Nuclear Ribosomal DNA. *Korean J. Parasitology*. **52**: 701-705.
- Mehlhorn, H. (2008). *Encyclopedia of Parasitology*. Springer Publication. 1461.
- Mohanta, U.K., Rana, H.B., Devkota, B. and Itagaki, T. (2017). Molecular and phylogenetic analyses of the liver amphistome *Explanatum explanatum* (Creplin, 1847) Fukui, 1929 in ruminants from Bangladesh and Nepal based on nuclear ribosomal ITS2 and mitochondrial nad1 sequences. *J. Helminthology*. **91**: 497-503.
- Prasad, A. and Varma, T.K. (1999). On the prevalence and community dominance among paramphistomes infecting domestic ruminants. *Veterinary Parasitology*. **13**: 129-133.
- Sambrook, J. and Russell, D.W. (2006). Purification of nucleic acids by extraction with phenol: chloroform. *CSH Protoc. pdb. prot.* 4455.
- Sanguankiat, S., Sato, M.O., Sato, M., Maipanich, W., Yoonuan, T., Pangvongsa, T., Boupah, B., Chigusa, Y., Moji, K. and Waikagul, J. (2016). First Record of Paramphistomes *Fiscoederius cobboldi* and *Paramphistomum epiclutum* Detected in Bovine Rumen from a Local Market of Savannakhet Province, Lao PDR. *Korean J. Parasitology*. **54(4)**: 543-547.
- Say, O. (1990). *CRC Handbook of the Zoology of Amphistomes*. CRC press Inc. Boca Ratón, Florida.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A. and Kumar, S. (2013). MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution*. **30**: 2725-2729.
- Vandna, B. and Gupta, R. (2019). Occurance of paramphistomosis in cattle and buffaloes with digestive disorders in ambala district of Haryana. *Haryana Vet*. **58(2)**: 298-300.
- Yamaguti, S. (1971). *Synopsis of digenetic trematodes of vertebrates*. Vols I and II. Keigaku Publishing Co. Tokyo, Japan. p. 1074.

THE HARYANA VETERINARIAN

Editors/Editorial Board Members are highly thankful to all the distinguished referees who helped us in the evaluation of articles. We request them to continue to extend their co-operation and be prompt in future to give their valuable comments on the articles for timely publication of the journal.